Emission Inventory Conference San Diego, California April 28-May 1, 2003



Emission Inventory Preparation for Air Quality Modelling in the Pacific Northwest

J. Wayne Boulton, Martin Gauthier, Xin Qiu and Mike Lepage RWDI West Inc. (RWDI Group of Companies)
Guelph, Ontario, Canada

Colin di Cenzo

Environment Canada, Pacific & Yukon Region Vancouver, British Columbia, Canada

PNW Modelling Project



Study design

- Prepare MC2 meteorological data for input to SMOKE and CMAQ
- Compile and process EI data through SMOKE
- Perform base case air quality modelling over
 12-km and 4-km domains for two episodes
 - Summer: August 9 to 20, 2001
 - Winter: December 1 to 12, 2002
- Perform air quality modelling for transboundary and future year emission scenarios


PNW Model Domains





MC2 ® MCIP2 Conversion

Environment Canada's MC2 model output used to run SMOKE and CMAQ

- Map model specific meteorological parameters
- Projection conversion
 (Polar Stereographic to Lambert Conic Conformal)
- Horizontal interpolation
- Vertical interpolation (sigma level computations)







Surface Wind and Temperature Fields

August 2001 Base Case. 12-km Grid Domain Pacific Northwest IAQMP True North

Video #2 Prepared by: VCT Project #W03-121A RWDI March 10, 2003

RWDI

Emission Inventory Data

- GVRD/FVRD, 2000 (GVRD)
- Whatcom county, year (GVRD)
- Canada/US marine, 2000 (GVRD)
- BC and Alberta, 1995 CAC EI (EC)
- US, 1996 mobile grown to 2003 (WSU)-
- US, 1996 area grown to 2001 (EPA / RWDI)
- Washington state, points, 1999 (Wash. DOE)
- Rest of US, 1996 point inventory (EPA)
- Biogenics processed through BEIS2 (RWDI)

RWDI's "SMOKE-In" EI Tool


RWDI

📠 SMOKE-in area : Database

Objects


gryUserExportTableCreate_GrowthFactor: Make Table

- Microsoft ACCESS Database Tool
- Import EC NET and SMOKE IDA files
- Combine multiple EI data Files
- Query, sort, growth factors, QA/QC, etc.
- **Export SMOKE IDA data files**




Allocation of Marine Emissions






Mobile Emissions: NOx

2001 BASE 4 km grid





RWDI

Mobile NOx Emissions


August 2001 Base Case. 4-km Grid Domain Pacific Northwest IAQMP




Video #1 Prepared by: VCT Project #W03-121A




O3






August 14,2001 0:00:00 Min= 16.3 at (27,47), Max= 76.8 at (18,49



August 2001 Base Case. 12-km Grid Domain Pacific Northwest IAQMP




Video #3 Prepared by: VCT Project #W03-121A






PM2.5

BASE 12 km grid



August 11,2001 14:00:00 Min= 0.3 at (6,2), Max=135.9 at (64,86)




PM 2.5 Concentration

August 2001 Base Case. 12-km Grid Domain Pacific Northwest IAQMP



Video #5 Prepared by: VCT Project #W03-121A






Vancouver


RWDI



Legend

ModelledMeasured










RWDI

Legend

ModelledMeasured



PNW Modelling Project



Technical Innovations by RWDI

- CMAQ recompiled and run on Linux in parallel on a 6-node cluster
- Developed software to convert MC2 to MM5
- Used 3.3-km MC2 data for 12-km and 4-km SMOKE and CMAQ runs
- Used SMOKE-In to prepare emission data
- Created software to crop, re-index, and interpolate gridded spatial surrogates
- Ran SMOKE v1.3 on Windows platform

PNW Modelling Project



Some Key Conclusions / Summary to Date

- Met data conversion successful
- Emission consolidation and processing successful (VERY time consuming)
- CMAQ results compare well with observations for ozone and PM_{2.5} at 12-km and 4-km res.
 - Daytime peaks and nighttime lows in ozone levels reproduced well in urban areas but overpredicted at night in rural areas
 - PM_{2.5} results better on 24-hour average